

**Sniffing COVID-19 Out: Studying the Nose to Understand the Coronavirus
Neuroscience Quarterly, Fall 2020**

References:

- Parma, V., Et al. (2020). More than smell. COVID-19 is associated with severe impairment of smell, taste, and chemesthesis. *Chemical Senses*. doi: [10.1093/chemse/bjaa041](https://doi.org/10.1093/chemse/bjaa041)
- Dubé, M., Le Coupanec, A., Wong, A. H. M., Rini, J. M., Desforges, M., Talbot, P. J. (2018). Axonal Transport Enables Neuron-to-Neuron Propagation of Human Coronavirus OC43. *Journal of Virology*, 92(17), e00404-18. doi: [10.1128/JVI.00404-18](https://doi.org/10.1128/JVI.00404-18)
- Durrant, D. M., Ghosh, S., & Klein R. S., (2016). The Olfactory Bulb: An Immunosensory Effector Organ during Neurotropic Viral Infections. *ACS Chemical Neuroscience*, 7(4): 464–469. doi:[10.1021/acscchemneuro.6b00043](https://doi.org/10.1021/acscchemneuro.6b00043)
- Cooper K. W., Et al. (2020). COVID-19 and the Chemical Senses: Supporting Players Take Center Stage. *Neuron*. doi: [10.1016/j.neuron.2020.06.032](https://doi.org/10.1016/j.neuron.2020.06.032)
- Menni C., Et al. (2020). Real-time tracking of self-reported symptoms to predict potential COVID-19. *Nature Medicine*, 26, 1037–1040. doi: 10.1038/s41591-020-0916-2
- Sun, S., Et al. (2020). A Mouse Model of SARS-CoV-2 Infection and Pathogenesis. *Cell Host & Microbe*, 28(1) 124-133.e4. doi: [10.1016/j.chom.2020.05.020](https://doi.org/10.1016/j.chom.2020.05.020)
- Brann, D. H., Et al. (2020). Non-neuronal expression of SARS-CoV-2 entry genes in the olfactory system suggests mechanisms underlying COVID-19-associated anosmia. BioRxiv. Doi: [10.1101/2020.03.25.009084](https://doi.org/10.1101/2020.03.25.009084)